INDEX TO AUTHORS

ABERNATHY, CHARLES. See FOLKMAN, MERLER, ABERNATHY, and WILLIAMS, 275

ACKERMAN, ALEKSANDER. See EIDINGER and ACKERMAN, 1061

ADLER, W. H. See TAKIGUCHI, ADLER, and SMITH, 63

AISENBERG, ALAN C., and MURRAY, CAROLINE. The contrasting effects of cyclophosphamide and radiation on the immune responses of the mouse, 877

AMOS, D. B. See SEIGLER, WARD, AMOS, PHAUP, and STICKEL, 411

ANDERSON, BYRON. See FEIZI, KABAT, VICARI, ANDERSON, and MARSH, 39

AOKI, TADAO. See GIRALDO, BETH, HIRSHAUT, AOKI, OLD, BOYSE, and CHOPRA, 454

—. See MELLORS, SHIRAI, AOKI, HUEBNER, and KRAWCZYNSKI, 113

AREND, WILLIAM P. See MANNIK, AREND, HALL, and GILLILAND, 713

AUSTEN, K. FRANK. See BACH, BLOCH, and AUSTEN, 752, 772

—. See KAPLAN and AUSTEN, 696

—. See KAY, STECHSCHULTE, and AUSTEN, 602

AXELRAD, MICHAEL A. See RADCLIFFE and AXELRAD, 846

—. Suppression of memory by passive immunization late in the primary response, 857

BACH, MICHAEL K., BLOCH, KURT J., and AUSTEN, K. FRANK. IgE and IgGa antibody-mediated release of histamine from rat peritoneal cells. I. Optimum conditions for in vitro preparation of target cells with antibody and challenge with antigen, 752

BACH, MICHAEL K., BLOCH, KURT J., and AUSTEN, K. FRANK. II. Interaction of IgGa and IgE at the target cell, 772

BARKER, CLYDE F., and BILLINGHAM, R. E. The lymphatic status of hamster cheek pouch tissue in relation to its properties as a graft and as a graft site, 620

BAST, ROBERT C., JR., MANSEAU, ELEANOR J., and DVORAK, HAROLD F. Heterogeneity of the cellular immune response. I. Kinetics of lymphocyte stimulation during sensitization and recovery from tolerance, 187

—, SIMPSON, BLANCHE A., and —. II. The role of adjuvant. Lymphocyte stimulation in cutaneous basophil hypersensitivity, 202

BENACERRAF, BARUJ. See KATZ, PAUL, GOIDL, and BENACERRAF, 169

BERNHEIMER, ALAN W. See FISCHETTI, GOTSCHLICH, and BERNHEIMER, 1105

BETH, ELKE. See GIRALDO, BETH, HIRSHAUT, AOKI, OLD, BOYSE, and CHOPRA, 454

BEY, ELKE, and HARINGTON, J. S. Cytotoxic effects of some mineral dusts on Syrian hamster peritoneal macrophages, 1149

BILLINGHAM, R. E. See BARKER and BILLINGHAM, 620

BLANDEN, R. V. Mechanisms of recovery from a generalized viral

infection: mousepox. II. Passive transfer of recovery mechanisms with immune lymphoid cells, 1074

BLANDEN, R. V. III. Regression of infectious foci, 1090

BLOCH, KURT J. See BACH, BLOCH, and AUSTEN, 752, 772

BLOOM, BARRY R. See JIMENEZ, BLOOM, BLUME, and OETTGEN, 740

BLOTH, BJÖRN, and SVEHAG, SVEN-ERIC. Further studies on the ultrastructure of dimeric IgA of human origin, 1035

BLUME, MYRON R. See JIMENEZ, BLOOM, BLUME, and OETTGEN, 740

BOND, JOHN H., JR., ENGEL, ROLF R., and LEVITT, MICHAEL D. Factors influencing pulmonary methane excretion in man. An indirect method of studying the *in situ* metabolism of the methane-producing colonic bacteria, 572

BORNSTEIN, PAUL. See LINDSLEY, MANNIK, and BORNSTEIN, 1309

BOYD, ARTHUR D. See RAPAPORT, BOYD, SPENCER, LOWER, DAUSSET, CANNON, and FERREBEE, 260

BOYSE, EDWARD A. See GIRALDO, BETH, HIRSHAUT, AOKI, OLD, BOYSE, and CHOPRA, 454

—. See STOCKERT, OLD, and BOYSE, 1334

BRINSTER, R. L. See PALM, HEYNER, and BRINSTER, 1282

BRUDERER, R. See HALBERT, BRUDERER, and LIN, 677

BUXBAUM, J., ZOLLA, SUSAN, SCHARFF, M. D., and FRANKLIN, E. C. Synthesis and assembly of immunoglobulins by malignant human plasmacytes and lymphocytes. II. Heterogeneity of assembly in cells producing IgM proteins, 1118

CAMPBELL, PRISCILLA. See UNANUE, GREY, RABELLINO, CAMPBELL, and SCHMITKE, 1188

CANNON, FLORENCE D. See RAPAPORT, BOYD, SPENCER, LOWER, DAUSSET, CANNON, and FERREBEE, 260

CHOPRA, HARISH C. See GIRALDO, BETH, HIRSHAUT, AOKI, OLD, BOYSE, and CHOPRA, 454,

CINADER, B., JEEJEEBHOY, H. F., KOH, S. W., and RABBAT, A. G. Immunosuppressive and graft-rejecting antibodies in heterologous anti-lymphocytic serum, 81

CLAMAN, HENRY N. See COHEN and CLAMAN, 1026

COCHRANE, C. G. See HENSON and COCHRANE, 554

COHEN, IRUN R., GLOBERSON, AMIELA, and FELDMAN, MICHAEL. Autosensitization in vitro, 834

—, —, and —. Rejection of tumor allografts by mouse spleen cells sensitized in vitro, 821

COHEN, J. JOHN, and CLAMAN, HENRY N. Thymus–marrow immunocompetence. V. Hydrocortisone-resistant cells and processes in the hemolytic antibody response of mice, 1026

COHEN, STANLEY, and WARD, PETER A. In vitro and in vivo activity of a lymphocyte and immune complex-dependent chemotactic factor for eosinophils, 133

COHN, MELVIN. See GREY, HIRST, and COHN, 289

COHN, ZANVIL. See GORDON and COHN, 321

COLON, SONIA. See RABELLINO, COLON, GREY, and UNANUE, 156

CONE, ROBERT E., and JOHNSON, ARTHUR G. Regulation of the immune system by synthetic polynucleotides. III. Action on antigen-reactive cells of thymic origin, 665

COOK, MARGERY L. See STEVENS and COOK, 19

COSTELLO, RICHARD, and IZUMI, TAKATERU. Measurement of resistance to experimental tuberculosis in albino mice. The immune phase, 362
—. See IZUMI and COSTELLO, 376
CUDKOWICZ, GUSTAVO. See MILLER and CUDKOWICZ, 973

DAUSSET, JEAN. See RAPAPORT, BOYD, SPENCER, LOWER, DAUSSET, CANNON, and FERREBEE, 260
DAVID, JOHN R. See NATHAN, KARNOVSKY, and DAVID, 1356
DUBOS, RENÉ. See LEE and DUBOS, 147
—. See LEE, GORDON, LEE, and DUBOS, 339
DVORAK, HAROLD F. See BAST, MANSEAU, and DVORAK, 187
—. See BAST, SIMPSON, and DVORAK, 202

EICHMANN, K., and GREENBLATT, J. Relationships between relative binding affinity and electrophoretic behavior of rabbit antibodies to streptococcal carbohydrates, 424
EIDINGER, DAVID, and ACKERMAN, ALEKSANDER. A cellular deficit in the reconstitutive capacity of immune populations of lymphoid cells demonstrable in studies of delayed hypersensitivity in mice. Evidence for thymus–bone marrow cell synergism, 1061
—. See KERBEL and EIDINGER, 1043
ELLIOTT, S. D., HAYWARD, JOHN, and LIU, T. Y. The presence of a Group A variant-like antigen in streptococci of other groups with special reference to Group N, 479
ENGEL, ROLF R. See BOND, ENGEL, and LEVITT, 572

FAULK, W. PAGE. See MCCORMICK, FAULK, FOX, and FUDENBERG, 1
FEIZI, TEN, KABAT, ELVIN A., VICARI, GIUSEPPE, ANDERSON, BYRON, MARSH, W. LAURENCE. Immunochemical studies on blood groups. XLVII. The I antigen complex–precursors in the A, B, H, Le^a, and Le^b blood group system–hemagglutination-inhibition studies, 39
FELDMAN, JOSEPH D. See POLLEY, MÜLLER-EBERHARD, and FELDMAN, 53
FELDMAN, MICHAEL. See COHEN, GLOBERSON, and FELDMAN, 821, 834
FERREBEE, JOSEPH W. See RAPAPORT, BOYD, SPENCER, LOWER, DAUSSET, CANNON, and FERREBEE, 260
FERRIS, BARBARA. See NACHMAN, FERRIS, and HIRSCH, 785, 807
FINKIELMAN, SAMUEL. See FISCHER-FERRARO, NAHMOD, GOLDSTEIN, and FINKIELMAN, 353
FISCHER-FERRARO, CATALINA, NAHMOD, VÍCTOR E., GOLDSTEIN, DANIEL J., and FINKIELMAN, SAMUEL. Angiotensin and renin in rat and dog brain, 353
FISCHETTI, VINCENT A., GOTSCHLICH, EMIL C., and BERNHEIMER, ALAN W. Purification and physical properties of Group C streptococcal phage-associated lysin, 1105
FOLKMAN, JUDAH, MERLER, EZIO, ABERNATHY, CHARLES, and WILLIAMS, GRETCHEN. Isolation of a tumor factor responsible for angiogenesis, 275
FORD, WILLIAM L., and SIMONSEN, MORTEN. The factor of immunization in the rat. The effect of allogeneic immunization on graft-*versus*-host activity, 938
FOX, H. See MCCORMICK, FAULK, FOX, and FUDENBERG, 1
FRANKLIN, E. C. See BUXBAUM,

ZOLLA, SCHARFF, and FRANKLIN, 1118
FUDENBERG, H. H. See MCCORMICK, FAULK, FOX, and FUDENBERG, 1

GILLILAND, BRUCE C. See MANNIK, AREND, HALL, and GILLILAND, 713
GIRALDO, GAETANO, BETH, ELKE, HIRSHAUT, YASHAR, AOKI, TADAO, OLD, LLOYD J., BOYSE, EDWARD A., and CHOPRA, HARISH C. Human sarcomas in culture. Foci of altered cells and a common antigen; induction of foci and antigen in human fibroblast cultures by filtrates, 454
GLOBERSON, AMIELA. See COHEN, GLOBERSON, and FELDMAN, 821, 834
GOIDL, EDMOND A. See KATZ, PAUL, GOIDL, and BENACERRAF, 169
GOLDSTEIN, DANIEL J. See FISCHER-FERRARO, NAHMOD, GOLDSTEIN, and FINKIELMAN, 353
GOODMAN, JOEL W. See SENYK, WILLIAMS, NITECKI, and GOODMAN, 1294
GORDON, JAMES. See LEE, GORDON, LEE, and DUBOS, 339
GORDON, JULIUS. See TAN and GORDON, 520
GORDON, SAIMON, and COHN, ZANVIL. Macrophage–melanocyte heterokaryons. II. The activation of macrophage DNA synthesis. Studies with inhibitors of RNA synthesis, 321
GOTSCHLICH, EMIL C. See FISCHETTI, GOTSCHLICH, and BERNHEIMER, 1105
GOWANS, J. L. See TILNEY and GOWANS 951
GREENBLATT, J. See EICHMANN and GREENBLATT, 424
GREY, HOWARD M., HIRST, JUDITH WEGMAN, and COHN, MELVIN. A new mouse immunoglobulin: IgG3, 289
—. See RABELLINO, COLON, GREY, and UNANUE, 156
—. See UNANUE, GREY, RABELLINO, CAMPBELL, and SCHMITKE, 1188
GUDAT, FRED G., HARRIS, T. N., HARRIS, SUSANNA, and HUMMELER, KLAUS. Studies on antibody-producing cells. II. Appearance of ^{3}H-thymidine–labeled rosette-forming cells, 305

HALBERT, S. P., BRUDERER, R., and LIN, T. M. In vitro organization of dissociated rat cardiac cells into beating three-dimensional structures, 677
HALL, ANTHONY P. See MANNIK, AREND, HALL, and GILLILAND, 713
HARINGTON, J. S. See BEY and HARINGTON, 1149
HARRIS, SUSANNA. See GUDAT, HARRIS, HARRIS, and HUMMELER, 305
HARRIS, T. N. See GUDAT, HARRIS, HARRIS, and HUMMELER, 305
HARTLEY, JANET W. See PINCUS, HARTLEY, and ROWE, 1219
HARTMANN, KLAUS-ULRICH. Induction of a hemolysin response in vitro. II. Influence of the thymus-derived cells during the development of the antibody-producing cells, 1325
HAYWARD, JOHN. See ELLIOTT, HAYWARD, and LIU, 479
HAYWOOD, GERALD R., and MCKHANN, CHARLES F. Antigenic specificities on murine sarcoma cells. Reciprocal relationship between normal transplantation antigens (H-2) and tumor-specific immunogenicity, 1171
HEGGIE, ALFRED D. Pathogenesis of H-1 virus infection of embryonic

hamster bone in organ culture, 506
HENSON, P. M., and COCHRANE, C. G. Acute immune complex disease in rabbits. The role of complement and of a leukocyte-dependent release of vasoactive amines from platelets, 554
HEYNER, SUSAN. See PALM, HEYNER, and BRINSTER, 1282
HILL, JEFFREY H., and WARD, PETER A. The phlogistic role of C3 leukotactic fragments in myocardial infarcts of rats, 885
HIRSCH, JAMES G. See JONES and HIRSCH, 231
—. See NACHMAN, FERRIS, and HIRSCH, 785, 807
HIRSHAUT, YASHAR. See GIRALDO, BETH, HIRSHAUT, AOKI, OLD, BOYSE, and CHOPRA, 454
HIRST, JUDITH WEGMAN. See GREY, HIRST, and COHN, 289
HUEBNER, ROBERT J. See MELLORS, SHIRAI, AOKI, HUEBNER, and KRAWCZYNSKI, 113
HUMMELER, KLAUS. See GUDAT, HARRIS, HARRIS, and HUMMELER, 305

IZUMI, TAKATERU, and COSTELLO, RICHARD. Temporal development of resistance to pulmonary tuberculosis in Swiss albino mice, 376
—. See COSTELLO and IZUMI, 362

JEEJEEBHOY, H. F. See CINADER, JEEJEEBHOY, KOH, and RABBAT, 81
JESAITIS, MARGERIS A. See WELTZIEN and JESAITIS, 534
JIMINEZ, LUIS, BLOOM, BARRY R., BLUME, MYRON R., and OETTGEN, HERBERT F. On the number and nature of antigen-sensitive lymphocytes in the blood of delayed-hypersensitive human donors, 740
JOHNSON, ARTHUR G. See CONE and JOHNSON, 665
—. See JOHNSON and JOHNSON, 649
JOHNSON, HERBERT G., and JOHNSON, ARTHUR G. Regulation of the immune system by synthetic polynucleotides. II. Action on peritoneal exudate cells, 649
JONES, THOMAS C., and HIRSCH, JAMES G. The interaction in vitro of *Mycoplasma pulmonis* with mouse peritoneal macrophages and L-cells, 231

KABAT, ELVIN A. See FEIZI, KABAT, VICARI, ANDERSON, and MARSH, 39
KAPLAN, ALLEN P., and AUSTEN, K. FRANK. A prealbumin activator of prekallikrein. II. Derivation of activators of prekallikrein from active Hageman factor by digestion with plasmin, 696
KARNOVSKY, MANFRED L. See NATHAN, KARNOVSKY, and DAVID, 1356
KATZ, DAVID H., PAUL, WILLIAM E., GOIDL, EDMOND A., and BENACERRAF, BARUJ. Carrier function in anti-hapten antibody responses. III. Stimulation of antibody synthesis and facilitation of hapten-specific secondary antibody responses by graft-*versus*-host reactions, 169
KAY, A. B., STECHSCHULTE, DANIEL J., and AUSTEN, K. FRANK. An eosinophil leukocyte chemotactic factor of anaphylaxis, 602
KERBEL, R. S., and EIDINGER, DAVID. Further studies of antigenic competition. III. A model to account for the phenomenon based on a deficiency of cell-to-cell interaction in immune lymphoid cell populations, 1043
KHAITOV, R. M. See PETROV, MANYKO, KHAITOV, and SESLAVINA, 640

KLINMAN, NORMAN R. The secondary immune response to a hapten in vitro. Antigen concentration and the carrier effect, 963
KOH, S. W. See CINADER, JEEJEEBHOY, KOH, and RABBAT, 81
KOSTER, F. T., and MCGREGOR, D. D. The mediator of cellular immunity. III. Lymphocyte traffic from the blood into the inflamed peritoneal cavity, 864
—, —, and MACKANESS, G. B. II. Migration of immunologically committed lymphocytes into inflammatory exudates, 400
—. See MCGREGOR, KOSTER, and MACKANESS, 389
KRAWCZYNSKI, KRZYSZTOF. See MELLORS, SHIRAI, AOKI, HUEBNER, and KRAWCZYNSKI, 113
KUNKEL, H. G. See NATVIG, MICHAELSEN, and KUNKEL, 1004

LAMM, MICHAEL E. See PINCUS, LAMM, and NUSSENZWEIG, 987
LEE, ADRIAN, GORDON, JAMES, LEE, CHI-JEN, and DUBOS, RENÉ. The mouse intestinal microflora with emphasis on the strict anaerobes, 339
LEE, CHI-JEN, and DUBOS, RENÉ. Lasting biological effects of early environmental influences. VI. Effects of early environmental stresses on metabolic activity and organ weights, 147
—. See LEE, GORDON, LEE, and DUBOS, 339
LEVIN, HOWARD A., LEVINE, HERBERT, and SCHLOSSMAN, STUART F. Antigen recognition and antibody specificity. Carrier specificity and genetic control of anti-dinitrophenyl-oligolysine antibody, 1199
LEVINE, BERNARD B. See PHILLIPS-QUAGLIATA, LEVINE, QUAGLIATA, and UHR, 589
LEVINE, HERBERT. See LEVIN, LEVINE, and SCHLOSSMAN, 1199
LEVITT, MICHAEL D. See BOND, ENGEL, and LEVITT, 572
LILLY, FRANK. See PINCUS, ROWE, and LILLY, 1234
LIN, T. M. See HALBERT, BRUDERER, and LIN, 677
LINDSLEY, HERBERT, MANNIK, MART, and BORNSTEIN, PAUL. The distribution of antigenic determinants in rat skin collagen, 1309
LIU, T. Y. See ELLIOTT, HAYWARD, and LIU, 479
LOWER, RICHARD R. See RAPAPORT, BOYD, SPENCER, LOWER, DAUSSET, CANNON, and FERREBEE, 260

MACKANESS, G. B. See KOSTER, MCGREGOR, and MACKANESS, 400
—. See MCGREGOR, KOSTER, and MACKANESS, 389
MANNIK, MART, AREND, WILLIAM P., HALL, ANTHONY P., and GILLILAND, BRUCE C. Studies on antigen–antibody complexes. I. Elimination of soluble complexes from rabbit circulation, 713
—. See LINDSLEY, MANNIK, and BORNSTEIN, 1309
MANSEAU, ELEANOR J. See BAST, MANSEAU, and DVORAK, 187
MANYKO, V. M. See PETROV, MANYKO, KHAITOV, and SESLAVINA, 640
MARSH, W. LAURENCE. See FEIZI, KABAT, VICARI, ANDERSON, and MARSH, 39
MARVER, HARVEY S. See PIMSTONE, TENHUNEN, SEITZ, MARVER, and SCHMID, 1264
MCCARTY, DANIEL J. See WALLINGFORD and MCCARTY, 100
MCCLUSKEY, ROBERT T. See WERDELIN and MCCLUSKEY, 1242
MCCORMICK, J. N., FAULK, W. PAGE, FOX, H., and FUDENBERG, H. H. Immunohistological and elution studies of the human placenta, 1

McGregor, D. D., Koster, F. T., and Mackaness, G. B. The mediator of cellular immunity. I. The life-span and circulation dynamics of the immunologically committed lymphocyte, 389
—. See Koster and McGregor, 864
—. See Koster, McGregor, and Mackaness, 400
McKhann, Charles F. See Haywood and McKhann, 1171
Mellors, Robert C., Shirai, Toshikazu, Aoki, Tadao, Huebner, Robert J., and Krawczynski, Krzysztof. Wild-type Gross leukemia virus and the pathogenesis of the glomerulonephritis of New Zealand mice, 113
Merler, Ezio. See Folkman, Merler, Abernathy, and Williams, 275
Michaelsen, T. E. See Natvig, Michaelsen, and Kunkel, 1004
Miller, Harold C., and Cudkowicz, Gustavo. Antigen-specific cells in mouse bone marrow. II. Fluctuation of the number and potential of immunocyte precursors after immunization, 973
Mozes, Edna. See Shearer, Mozes, and Sela, 216
Müller-Eberhard, Hans J. See Polley, Müller-Eberhard, and Feldman, 53
Murray, Caroline. See Aisenberg and Murray, 877

Nachman, Ralph L., Ferris, Barbara, and Hirsch, James G. Macrophage plasma membranes. I. Isolation and studies on protein components, 785
—, —, and —. II. Studies on synthesis and turnover of protein constituents, 807
Nahmod, Víctor E. See Fischer-Ferraro, Nahmod, Goldstein, and Finkielman, 353
Nathan, Carl F., Karnovsky, Manfred L., and David, John R. Alterations of macrophage functions by mediators from lymphocytes, 1356
Natvig, J. B., Michaelsen, T. E., and Kunkel, H. G. Evidence for recent duplications among certain γ-globulin heavy chain genes, 1004
Nitecki, Danute E. See Senyk, Williams, Nitecki, and Goodman, 1294
Nowell, Peter C., and Wilson, Darcy B. Studies on the life history of lymphocytes. I. The life-span of cells responsive in the mixed lymphocyte interaction, 1131
—. See Wilson and Nowell, 442
Nussenzweig, Victor. See Pincus, Lamm, and Nussenzweig, 987

Oettgen, Herbert F. See Jimenez, Bloom, Blume, and Oettgen, 740
Old, Lloyd J. See Giraldo, Beth, Hirshaut, Aoki, Old, Boyse, and Chopra, 454
—. See Stockert, Old, and Boyse, 1334

Palm, Joy, Heyner, Susan, and Brinster, R. L. Differential immunofluorescence of fertilized mouse eggs with *H-2* and non–*H-2* antibody, 1282
Paul, William E. See Katz, Paul, Goidl, and Benacerraf, 169
Pearson, Margot N., and Raffel, Sidney. Macrophage-digested antigen as inducer of delayed hypersensitivity, 494
Petrov, R. V., Manyko, V. M., Khaitov, R. M., and Seslavina, L. S. An experimental system for the simultaneous estimation of mitostatic and lymphotoxic effects of immunosuppressants and cytostatics, 640

PHAUP, M. B. See SEIGLER, WARD, AMOS, PHAUP, and STICKEL, 411

PHILLIPS-QUAGLIATA, JULIA M., LEVINE, BERNARD B., QUAGLIATA, FRANCO, and UHR, JONATHAN W. Mechanisms underlying binding of immune complexes to macrophages, 589

PIMSTONE, NEVILLE R., TENHUNEN, RAIMO, SEITZ, PAUL T., MARVER, HARVEY S., and SCHMID, RUDI. The enzymatic degradation of hemoglobin to bile pigments by macrophages, 1264

PINCUS, CAROLYN S., LAMM, MICHAEL E., and NUSSENZWEIG, VICTOR. Regulation of the immune response: suppressive and enhancing effects of passively administered antibody, 987

PINCUS, THEODORE, HARTLEY, JANET W., and ROWE, WALLACE P. A major genetic locus affecting resistance to infection with murine leukemia viruses. I. Tissue culture studies of naturally occurring viruses, 1219

—, ROWE, WALLACE P., and LILLY, FRANK. II. Apparent identity to a major locus described for resistance to Friend murine leukemia virus, 1234

POLLEY, MARGARET J., MÜLLER-EBERHARD, HANS J., and FELDMAN, JOSEPH D. Production of ultrastructural membrane lesions by the fifth component of complement, 53

QUAGLIATA, FRANCO. See PHILLIPS-QUAGLIATA, LEVINE, QUAGLIATA, and UHR, 589

RABBAT, A. G. See CINDER, JEEJEEBHOY, KOH, and RABBAT, 81

RABELLINO, ENRIQUE, COLON, SONIA, GREY, HOWARD M., and UNANUE, EMIL R. Immunoglobulins on the surface of lymphocytes. I. Distribution and quantitation, 156

—. See UNANUE, GREY, RABELLINO, CAMPBELL, and SCHMITKE, 1188

RADCLIFFE, GORDON N., and AXELRAD, MICHAEL A. Immunological memory in vitro, 846

RAFFEL, SIDNEY. See PEARSON and RAFFEL, 494

RAPAPORT, FELIX T., BOYD, ARTHUR D., SPENCER, FRANK C., LOWER, RICHARD R., DAUSSET, JEAN, CANNON, FLORENCE D., and FERREBEE, JOSEPH W. Histocompatibility studies in a closely bred colony of dogs. II. Influence of the DL-A system of canine histocompatibility upon the survival of cardiac allografts, 260

ROWE, WALLACE P. See PINCUS, HARTLEY, and ROWE, 1219

—. See PINCUS, ROWE, and LILLY, 1234

SCHARFF, M. D. See BUXBAUM, ZOLLA, SCHARFF, and FRANKLIN, 1118

SCHLOSSMAN, STUART F. See LEVIN, LEVINE, and SCHLOSSMAN, 1199

SCHMID, RUDI. See PIMSTONE, TENHUNEN, SEITZ, MARVER, and SCHMID, 1264

SCHMITKE, JON. See UNANUE, GREY, RABELLINO, CAMPBELL, and SCHMITKE, 1188

SEIGLER, H. F., WARD, F. E., AMOS, D. B., PHAUP, M. B., and STICKEL, D. L. The immunogenicity of human HL-A haplotypes as measured by skin graft survival times and mixed leukocyte reactions, 411

SEITZ, PAUL T. See PIMSTONE, TENHUNEN, SEITZ, MARVER, and SCHMID, 1264

SELA, MICHAEL. See SHEARER, MOZES, and SELA, 216
SENYK, GEORGE, WILLIAMS, E. BRADY, NITECKI, DANUTE E., and GOODMAN, JOEL W. The functional dissection of an antigen molecule: specificity of humoral and cellular immune responses to glucagon, 1294
SESLAVINA, L. S. See PETROV, MANYKO, KHAITOV, and SESLAVINA, 640
SHEAGREN, JOHN N. See SIMON and SHEAGREN, 1377
SHEARER, G. M., MOZES, EDNA, and SELA, MICHAEL. Cellular basis of the genetic control of immune responses to synthetic polypeptides. II. Frequency of immunocompetent precursors specific for two distinct regions within (Phe, G)–Pro–-L, a synthetic polypeptide derived from multichain polyproline, in inbred mouse strains, 216
SHERR, CHARLES J., and UHR, JONATHAN W. Immunoglobulin synthesis and secretion. VI. Synthesis and intracellular transport of immunoglobulin in nonsecretory lymphoma cells, 901
SHIRAI, TOSHIKAZU. See MELLORS, SHIRAI, AOKI, HUEBNER, and KRAWCZYNSKI, 113
SILVERS, WILLYS K. See WACHTEL and SILVERS, 921
SIMON, HARVEY B., and SHEAGREN, JOHN N. Cellular immunity in vitro. I. Immunologically mediated enhancement of macrophage bactericidal capacity, 1377
SIMONSEN, MORTEN. See FORD and SIMONSEN, 938
SIMPSON, BLANCHE A. See BAST, SIMPSON, and DVORAK, 202
SJÖBERG, OLOF. Antigen-binding cells in mice immune or tolerant to *Escherichia coli* polysaccharide, 1015
SMITH, R. T. See TAKIGUCHI, ADLER, and SMITH, 63
SPENCER, FRANK C. See RAPAPORT, BOYD, SPENCER, LOWER, DAUSSET, CANNON, and FERREBEE, 260
STECHSCHULTE, DANIEL J. See KAY, STECHSCHULTE, and AUSTEN, 602
STEVENS, JACK G., and COOK, MARGERY L. Restriction of herpes simplex virus by macrophages. An analysis of the cell-virus infection, 19
STICKEL, D. L. See SEIGLER, WARD, AMOS, PHAUP, and STICKEL, 411
STOCKERT, ELISABETH, OLD, LLOYD J., and BOYSE, EDWARD A. The G_{IX} system. A cell surface alloantigen associated with murine leukemia virus; implications regarding chromosomal integration of the viral genome, 1334
SVEHAG, SVEN-ERIC. See BLOTH and SVEHAG. 1035

TAKIGUCHI, T., ADLER, W. H., and SMITH, R. T. Cellular recognition in vitro by mouse lymphocytes. Effects of neonatal thymectomy and thymus graft restoration on alloantigen and PHA stimulation of whole and gradient-separated subpopulations of spleen cells, 63
TAN, TERESITA, and GORDON, JULIUS. Participation of three cell types in the anti-sheep red blood cell response in vitro. Separation of antigen-reactive cells from the precursors of antibody-forming cells, 520
TENHUNEN, RAIMO. See PIMSTONE, TENHUNEN, SEITZ, MARVER, and SCHMID, 1264
TILNEY, N. L., and GOWANS, J. L. The sensitization of rats by allografts transplanted to alymphatic pedicles of skin, 951

UHR, JOHATHAN W. See PHILLIPS-QUAGLIATA, LEVINE, QUAGLIATA, and UHR, 589
—. See SHERR and UHR, 901

UNANUE, EMIL R., GREY, HOWARD M., RABELLINO, ENRIQUE, CAMPBELL, PRISCILLA, and SCHMITKE, JON. Immunoglobulins on the surface of lymphocytes. II. The bone marrow as the main source of lymphocytes with detectable surface-bound immunoglobulin, 1188

—. See RABELLINO, COLON, GREY, and UNANUE, 156

VICARI, GIUSEPPE. See FEIZI, KABAT, VICARI, ANDERSON, and MARSH, 39

WACHTEL, STEPHEN S., and SILVERS, WILLYS K. Skin homografts: tolerogenic *versus* immunogenic influences in mice, 921

WALLINGFORD, WALTER R., and MCCARTY, DANIEL J. Differential membranolytic effects of microcrystalline sodium urate and calcium pyrophosphate dihydrate, 100

WARD, F. E. See SEIGLER, WARD, AMOS, PHAUP, and STICKEL, 411

WARD, PETER A. See COHEN and WARD, 133

—. See HILL and WARD, 885

WELTZIEN, HANS ULRICH, and JESAITIS, MARGERIS A. The nature of the colicin K receptor of *Escherichia coli* Cullen, 534

WERDELIN, OLE, and MCCLUSKEY, ROBERT T. The nature and the specificity of mononuclear cells in experimental autoimmune inflammations and the mechanisms leading to their accumulation, 1242

WILLIAMS, E. BRADY. See SENYK, WILLIAMS, NITECKI, and GOODMAN, 1294

WILLIAMS, GRETCHEN. See FOLKMAN, MERLER, ABERNATHY, and WILLIAMS, 275

WILSON, DARCY B., and NOWELL, PETER C. Quantiative studies on the mixed lymphocyte interaction in rats. V. Tempo and specificity of the proliferative response and the number of reactive cells from immunized donors, 442

—. See NOWELL and WILSON, 1131

ZOLLA, SUSAN. See BUXBAUM, ZOLLA, SCHARFF, and FRANKLIN, 1118

INDEX TO SUBJECTS

ADJUVANT, role of. Lymphocyte stimulation in cutaneous basophil hypersensitivity, 202

Alloantigen and PHA stimulation of whole and gradient-separated subpopulations of spleen cells, effects of neonatal thymectomy and thymus graft restoration on, 63

Allo-antigen, cell surface (G1x system) associated with murine leukemia virus; implications regarding chromosomal integration of viral genome, 1334

Allogeneic immunization, effect of, on graft-*versus*-host activity, 938

Allografts, cardiac, survival of, influence of DL-A system of canine histocompatibility upon, 260

—, transplanted to alymphatic pedicles of skin, sensitization of rats by, 951

—, tumor, rejection of, by mouse spleen cells sensitized in vitro, 821

Alymphatic pedicles of skin, allografts transplanted to, sensitzation of rats by, 951

Amines, vasoactive, leukocyte-dependent release of from platelets, and complement, role of, 554

Anaerobes, strict, mouse intestinal microflora with emphasis on, 339

Anaphylaxis, eosinophil leukocyte chemotactic factor of, 602

Angiogenesis, tumor factor responsible for, isolation of, 275

Angiotensin and renin in rat and dog brain, 353

Antibodies, immunosuppressive and graft-rejecting, in heterologous antilymphocytic serum, 81

Antibodies, rabbit, to streptococcal carbohydrates, relationship between relative binding affinity and electrophoretic behavior of, 424

Antibody, H-2 and non–H-2, differential immunofluorescence of fertilized mouse eggs with, 1282

—, hemolytic, response of mice, hydrocortisone-resistant cells and processes in, 1026

—, IgE and IgGa, -mediated release of histamine from rat peritoneal cells, 752, 772

—, in vitro preparation of target cells with, and challenge with antigen, optimum conditions for, 752

—, passively administered, suppressive and enhancing effects of: regulation of immune response, 987

— responses, hapten-specific secondary, facilitation of, and stimulation of antibody synthesis, by graft-*versus*-host reactions, 169

— specificity and antigen recognition. Carrier specificity and genetic control of anti–dinitrophenyl-oligolysine antibody, 1199

— synthesis, stimulation of, and facilitation of hapten-specific secondary antibody responses by graft-*versus*-host reactions, 169

Antibody-forming cells, precursors of, separation from antigen-reactive cells, 520

Antibody-producing cells, influence of thymus-derived cells during development of, 1325

Antigen, challenge with, and in vitro preparation of target cells with antibody, optimum conditions for, 752

Antigen complex, I,–precursors in A, B, H, Le^a, and Le^b blood group system–hemagglutination-inhibition studies, 39
— concentration and carrier effect: secondary immune response to hapten in vitro, 963
—, Group A variant-like, presence of in streptococci of other groups with special reference to Group N, 479
—, macrophage-digested, as inducer of delayed hypersensitivity, 494
— molecule, functional dissection of: specificity of humoral and cellular immune responses to glucagon, 1294
— recognition and antibody specificity. Carrier specificity and genetic control of anti–dinitrophenyl-oligolysine antibody, 1199
Antigen-binding cells in mice immune or tolerant to *Escherichia coli* polysaccharide, 1015
Antigen-reactive cells of thymic origin, action on, 665
— cells, separation of, from precursors of antibody-forming cells, 520
Antigen-sensitive lymphocytes, number and nature of, in blood of delayed-hypersensitive human donors, 740
Antigen-specific cells in mouse bone marrow, 973
Antigenic competition, further studies of. Model to account for phenomenon based on deficiency of cell-to-cell interaction in immune lymphoid cell populations, 1043
— determinants, distribution of, in rat skin collagen, 1309
— specificities on murine sarcoma cells. Reciprocal relationship between normal transplantation antigens (H-2) and tumor-specific immunogenicity, 1171
Antigens, normal transpantation (H-2) and tumor-specific immunogenicity, reciprocal relationship between. Antigenic specificities on murine sarcoma cells, 1171
Anti-hapten antibody responses, carrier function in, 169
Anti-lymphocytic serum, heterologous, immunosuppressive and graft-rejecting antibodies in, 81
Autosensitization in vitro, 834

BACTERIA, colonic, methane-producing, *in situ* metabolism of, indirect method of studying, 572
Basophil hypersensitivity, cutaneous, lymphocyte stimulation in. Role of adjuvant, 202
Bile pigments, enzymatic degradation of hemoglobin to, by macrophages, 1264
Binding affinity, relative, and electrophoretic behavior, relationships between, of rabbit antibodies to streptococcal carbohydrates, 424
— of immune complexes to macrophages, mechanisms underlying, 589
Blood group system, A, B, H, Le^a, and Le^b, precursors in–I antigen complex–hemagglutination-inhibition studies, 39
—, lymphocyte traffic from, into inflamed peritoneal cavity, 864
— of delayed-hypersensitive human donors, antigen-sensitive lymphocytes in, number and nature of, 740
Bone marrow as main source of lymphocytes with detectable surface-bound immunoglobulin, 1188
— marrow, mouse, antigen-specific cells in, 973
— marrow–thymus cell synergism, evidence for: cellular deficit in reconstitutive capacity of immune populations of lymphoid cells demonstrable in studies of delayed hypersensitivity in mice, 1061
Brain, rat and dog, angiotensin and renin in, 353

C3 leukotactic fragments, phlogistic role of, in myocardial infarcts of rats, 885
Calcium pyrophosphate dihydrate and microcrystalline sodium urate, differential membranolytic effects of, 100
Carbohydrates, streptococcal, rabbit antibodies to, relationship between relative binding affinity and electrophoretic behavior of, 424
Cardiac allografts, survival of, influence of DL-A system of canine histocompatibility upon, 260
— cells, rat, dissociated, in vitro organization of into beating three-dimensional structures, 677
Carrier effect and antigen concentration: secondary immune response to a hapten in vitro, 963
— function in anti-hapten antibody responses, 169
—specificity and genetic control of anti–dinitrophenyl-oligolysine antibody. Antigen recognition and antibody specificity, 1199
Cell populations, lymphoid, immune, cell-to-cell interaction in, model to account for phenomenon of antigenic competition based on deficiency of, 1043
—, red blood. *See* Erythrocyte.
—, target, interaction at, of IgGa and IgE, 772
Cell-to-cell interaction in immune lymphoid cell populations, model to account for phenomenon of antigenic competition based on deficiency of, 1043
Cell–virus interaction, analysis of, in macrophage restriction of herpes simplex virus, 19
Cells, altered, foci of, and a common antigen; induction of foci and antigen in human fibroblast cultures by filtrates, 454
— and processes, hydrocortisone-resistant, in hemolytic antibody response of mice, 1026
—, antibody-forming, precursors of, separation from antigen-reactive cells, 520
Cells, antibody-producing, influence of thymus-derived cells during development of, 1325
—, antigen-binding, in mice immune or tolerant to *Escherichia coli* polysaccharide, 1015
—, antigen-reactive, of thymic origin, action on, 665
—, antigen-reactive, separation of from precursors of antibody-forming cells, 520
—, antigen-specific, in mouse bone marrow, 973
—, L-, and mouse peritoneal macrophages, interaction in vitro of *Mycoplasma pulmonis* with, 231
—, lymphoid, immune populations of, reconstitutive capacity of, cellular deficit in, demonstrable in studies of delayed hypersensitivity in mice: evidence for thymus–bone marrow cell synergism, 1061
—, lymphoma, nonsecretory, immunoglobulin in, synthesis and intracellular transport of, 901
—, mononuclear, nature and specificity of, in experimental autoimmune inflammations and mechanisms leading to their accumulation, 1242
—, mouse spleen sensitized in vitro, rejection of tumor allografts by, 821
—, murine sarcoma, antigenic specificities on. Reciprocal relationship between normal transplantation antigens (H-2) and tumor-specific immunogenicity, 1171
—, peritoneal exudate, action on, 649
—, rat cardiac, dissociated, in vitro organization of into beating three-dimensional structures, 677
—, rat peritoneal, IgE and IgGa antibody-mediated release of histamine from, 752, 772
—, reactive, number of, and proliferative response from immunized

donors, tempo and specificity of, 442
Cells, spleen, whole and gradient-separated subpopulations of, effects of neonatal thymectomy and thymus graft restoration on alloantigen and PHA stimulation of, 63
—, target, in vitro preparation of with antibody and challenge with antigen, optimum conditions for, 752
—, thymus-derived, influence of during development of antibody-producing cells, 1325
Cellular basis of genetic control of immune responses to synthetic polypeptides, 216
— deficit in reconstitutive capacity of immune populations of lymphoid cells demonstrable in studies of delayed hypersensitivity in mice: evidence for thymus–bone marrow cell synergism, 1061
— immune and humoral responses to glucagon, specificity of: functional dissection of antigen molecule, 1294
— immune response, heterogeneity of, 187, 202
— immunity in vitro. Immunologically mediated enhancement of macrophage bactericidal capacity, 1377
— recognition in vitro by mouse lymphocytes, 63
Chemotactic factor, eosinophil leukocyte, of anaphylaxis, 602
— factor for eosinophils, immune complex-dependent, and lymphocyte, in vitro and in vivo activity of, 133
Chromosomal integration of viral genome, implications regarding; G_{1X} system: cell surface allo-antigen associated with murine leukemia virus, 1334
Circulation, rabbit, elimination of soluble (antigen–antibody) complexes from, 713
Collagen, rat skin, distribution of antigenic determinants in, 1309
Complement and leukocyte-dependent release of vasoactive amines from platelets, role of 554
—, fifth component of, production of ultrastructural membrane lesions by, 53
Complex disease, acute immune, in rabbits, 554
Complexes (antigen–antibody), soluble, elimination of, from rabbit circulation, 713
—, immune, binding of to macrophages, mechanisms underlying, 589
Colicin K receptor of *Escherichia coli* Cullen, nature of, 534
Cutaneous basophil hypersensitivity, lymphocyte stimulation in. Role of adjuvant, 202
Cytophosphamide and radiation, contrasting effects of, on immune responses of mouse, 877
Cytostatics and immunosuppressants, mitostatic and lymphotoxic effects of, experimental system for simultaneous estimation of, 640
Cytotoxic effects of mineral dusts on Syrian hamster peritoneal macrophages, 1149

DEOXYRIBONUCLEIC acid synthesis, macrophage, activation of. Studies with inhibitors of RNA synthesis, 321
Dihydrate, calcium pyrophosphate, and microcrystalline sodium urate, differential membranolytic effects of, 100
Dimeric IgA of human origin, ultrastructure of, further studies on, 1035
Dinitrophenyl, anti–, -oligolysine antibody, carrier specificity and genetic control of. Antigen recognition and antibody specificity, 1199
Dissection, functional, of antigen molecule: specificity of humoral and cellular immune responses to glucagon, 1294

DL-A system of canine histocompatibility, influence of, upon survival of cardiac allografts, 260

ELECTROPHORETIC behavior and relative binding affinity, relationships between, of rabbit antibodies to streptococcal carbohydrates, 424
Elution, and immunohistological studies of human placenta, 1
Embryonic hamster bone in organ culture, H-1 virus infection of, pathogenesis of, 506
Environmental stresses, early, effects of, on metabolic activity and organ weights, 147
Enzymatic degradation of hemoglobin to bile pigments by macrophage, 1264
Eosinophil leukocyte chemotactic factor of anaphylaxis, 602
Eosinophils, lymphocyte and immune complex-dependent chemotactic factor for, in vitro and in vivo activity of, 133
Erythrocyte, anti-sheep red blood cell response in vitro, participation of three cell types in, 520
Escherichia coli Cullen, colicin K receptor of, nature of, 534
— *coli* polysaccharide, antigen-binding cells in mice immune or tolerant to, 1015
Excretion, pulmonary methane, in man, factors influencing, 572
Exudates, inflammatory, migration of immunologically committed lymphocytes into, 400

FIBROBLAST cultures, human, induction of foci and antigen in by filtrates; foci of altered cells and a common antigen, 454
Filtrates, induction of foci and antigen in human fibroblast cultures by; foci of altered cells and a common antigen, 454
Foci and antigen, induction of, in human fibroblast cultures by filtrates; foci of altered cells and a common antigen, 454
— of altered cells and a common antigen; induction of foci and antigen in human fibroblast cultures by filtrates, 454
Friend murine leukemia virus, apparent identity to major locus described for resistance to, 1234

GIX system: cell surface allo-antigen associated with murine leukemia virus; implications regarding chromosomal integration of viral genome, 1334
γ-globulin heavy chain genes, evidence for recent duplications among, 1004
Genes, heavy chain, γ-globulin, evidence for recent duplications among, 1004
Genetic control of anti–dinitrophenyl-oligolysine antibody, and carrier specificity of. Antigen recognition and antibody specificity, 1199
— control of immune responses to synthetic polypeptides, cellular basis of, 216
— locus, major, affecting resistance to infection with murine leukemia viruses, 1219, 1234
Glomerulonephritis, pathogenesis of, and wild-type Gross leukemia virus, of New Zealand mice, 113
Glucagon, humoral and cellular immune responses to, specificity of: functional dissection of antigen molecule, 1294
Graft and graft site, lymphatic status of hamster cheek pouch tissue in relation to its properties as, 620
—, skin, survival times, and mixed leukocyte reactions, immunogenicity of human HL-A haplotypes as measured by, 411

Graft-*versus*-host activity, effect on, of allogeneic immunization, 938
— reactions, stimulation of antibody synthesis and facilitation of hapten-specific secondary antibody responses by, 169
Grafts, homo-, skin: tolerogenic vs. immunogenic influences in mice, 921
Gross leukemia virus, wild-type, and pathogenesis of glomerulonephritis of New Zealand mice, 113
Group C streptococcal phage-associated lysin, purification and physical properties of, 1105

H-1 virus infection of embryonic hamster bone in organ culture, pathogenesis of, 506
H-2 and non–H-2 antibody, differential immunofluorescence of fertilized mouse eggs with, 1282
Hageman factor, active, derivation of activators of prekallikrein from, by digestion with plasmin, 696
Haplotypes, human HL-A, immunogenicity of, as measured by skin graft survival times and mixed leukocyte reactions, 411
Hapten, anti-, antibody responses, carrier function in, 169
—, secondary immune response to in vitro: antigen concentration and carrier effect, 963
Hapten-specific secondary antibody responses, facilitation of, and stimulation of antibody synthesis, by graft-*versus*-host reactions, 169
Hemagglutination-inhibition studies–I antigen complex–precursors in A, B, H, Le^a, and Le^b blood group system, 39
Hemoglobin, enzymatic degradation of, to bile pigments by macrophages, 1264
Hemolysin response, induction of, in vitro. Influence of thymus-derived cells during development of antibody-producing cells, 1325
Hemolytic antibody response of mice, hydrocortisone-resistant cells and processes in, 1026
Herpes simplex virus, restriction of by macrophages, 19
Heterogeneity of cellular immune response, 187, 202
Heterokaryons, macrophage–melanocyte, 321
Histamine, IgE and IgGa antibody-mediated release of, from rat peritoneal cells, 752, 772
Histocompatibility, canine, influence of DL-A system of upon survival of cardiac allografts, 260
HL-A haplotypes, human, immunogenicity of, as measured by skin graft survival times and mixed leukocyte reactions, 411
Homografts, skin: tolerogenic vs. immunogenic influences in mice, 921
Human. *See* Man.
Humoral and cellular immune responses to glucagon, specificity of: functional dissection of antigen molecule, 1294
Hydrocortisone-resistant cells and processes in hemolytic antibody response of mice, 1026
Hypersensitivity, cutaneous basophil, lymphocyte stimulation in. Role of adjuvant, 202
—, delayed, in mice, cellular deficit in reconstitutive capacity of immune populations of lymphoid cells demonstrable in studies of: evidence for thymus–bone marrow cell synergism, 1061
—, delayed, macrophage-digested antigen as inducer of, 494

I ANTIGEN complex–precursors in A, B, H, Le^a, and Le^b blood group system–hemagglutination-inhibition studies, 39
IgA, dimeric, of human origin, ultrastructure of, further studies on, 1035

IgE and IgGa antibody-mediated release of histamine from rat peritoneal cells, 752, 772
— and IgGa, interaction of, at target cell, 772
IgGa and IgE antibody-mediated release of histamine from rat peritoneal cells, 752, 772
— and IgE, interaction of, at target cell, 772
IgG3, new mouse immunoglobulin, 289
Immune complex-dependent chemotactic factor for eosinophils, and lymphocyte, in vitro and in vivo activity of, 133
— complexes, binding of, to macrophages, mechanisms underlying, 589
— lymphoid cell populations, cell-to-cell interaction in, model to account for phenomenon of antigenic competition based on deficiency of, 1043
— populations of lymphoid cells, reconstitutive capacity of, cellular deficit in, demonstrable in studies of delayed hypersensitivity in mice: evidence for thymus–bone marrow cell synergism, 1061
— response, cellular, heterogeneity of, 187, 202
— response, regulation of: suppressive and enhancing effects of passively administered antibody, 987
— response, secondary, to hapten in vitro: antigen concentration and carrier effect, 963
— responses, cellular, and humoral responses to glucagon, specificity of: functional dissection of antigen molecule, 1294
— responses of mouse, contrasting effects on, of cyclophosphamide and radiation, 877
— responses to synthetic polypeptides, cellular basis of genetic control of, 216
— system, regulation of, by synthetic polynucleotides, 649, 665
Immunization, allogeneic, effect of, on graft-*versus*-host activity, 938
—, passive, suppression of memory by, late in primary response, 857
Immunized donors, proliferative response and number of reactive cells from, tempo and specificity of, 442
Immunochemical studies on blood groups, 39
Immunocompetent precursors, frequency of, specific for two distinct regions within (Phe, G)–Pro– –L, a synthetic polypeptide derived from multichain polyproline, in inbred mouse strains, 216
Immunofluorescence, differential, of fertilized mouse eggs with H-2 and non–H-2 antibody, 1282
Immunogenic vs. tolerogenic influences of skin homografts in mice, 921
Immunogenicity of human HL-A haplotypes as measured by skin graft survival times and mixed leukocyte reactions, 411
—, tumor-specific, and normal transplantation antigens (H-2), reciprocal relationship between. Antigenic specificities on murine sarcoma cells, 1171
Immunoglobulin, detectable surface-bound, lymphocytes with, bone marrow as main source of, 1188
—, new mouse: IgG3, 289
—, synthesis and intracellular transport of, in nonsecretory lymphoma cells, 901
Immunoglobulins on surface of lymphocytes, 156, 1188
—, synthesis and assembly of, by malignant human plasmacytes and lymphocytes, 1118
Immunohistological and elution studies of human placenta, 1
Immunological memory in vitro, 846

Immunologically committed lymphocyte, life-span and circulation dynamics of, 389
— committed lymphocytes, migration of, into inflammatory exudates, 400
— mediated enhancement of macrophage bactericidal capacity, 1377
Immunosuppressants and cytostatics, mitostatic and lymphotoxic effects of, experimental system for simultaneous estimation of, 640
Immunosuppressive and graft-rejecting antibodies in heterologous anti-lymphocytic serum, 81
Infection, resistance to, with murine leukemia viruses, major genetic locus affecting, 1219, 1234
—, viral, generalized: mousepox-mechanisms of recovery from, 1074, 1090
Inflammations, experimental autoimmune, nature and specificity of mononuclear cells in, and mechanisms leading to their accumulation, 1242
Inflammatory exudates, migration of immunologically committed lymphocytes into, 400
Intestinal microflora, mouse, with emphasis on strict anaerobes, 339
Intracellular transport and synthesis of immunoglobulin in nonsecretory lymphoma cells, 901

KALLIKREIN, pre-, activators of, derivation of from active Hageman factor by digestion with plasmin, 696
Kinetics of lymphocyte stimulation during sensitization and recovery from tolerance, 187

L-CELLS, and mouse peritoneal macrophages, interaction in vitro of *Mycoplasma pulmonis* with, 231
Lesions, membrane, ultrastructural, production of by fifth component of complement, 53
Leukemia virus, Gross, wild-type, and pathogensis of glomerulonephritis of New Zealand mice, 113
— virus, murine, cell surface alloantigen (G_{1X} system) associated with; implications regarding chromosomal integration of viral genome, 1334
— viruses, murine, major genetic locus affecting resistance to infection with, 1219, 1234
Leukocyte chemotactic factor, eosinophil, of anaphylaxis, 602
—, mixed, reactions, and skin graft survival times, immunogenicity of human HL-A haplotypes as measured by, 411
Leukocyte-dependent release of vasoactive amines from platelets, and complement, role of 554
Leukotactic fragments, C3, phlogistic role of, in myocardial infarcts of rats, 885
Life-span of cells responsive in mixed lymphocyte interaction, 1131
Lymphatic status of hamster cheek pouch tissue in relation to its properties as graft and graft site, 620
Lymphocyte and immune complex-dependent chemotactic factor for eosinophils, in vitro and in vivo activity of, 133
—, immunologically committed, life-span and circulation dynamics of, 389
— interaction, mixed, life-span of cells responsive in, 1131
— interaction, mixed, quantitative studies on, in rats, 442
— stimulation in cutaneous basophil hypersensitivity. Role of adjuvant, 202
— stimulation, kinetics of, during sensitization and recovery from tolerance, 187
— traffic from blood into inflamed peritoneal cavity, 864

Lymphocytes, alterations of macrophage functions by mediators from, 1356
—, and plasmacytes, malignant human, synthesis and assembly of immunoglobulins by, 1118
—, antigen-sensitive, number and nature of, in blood of delayed-hypersensitive human donors, 740
—, immunologically committed, migration of into inflammatory exudates, 400
—, mouse, cellular recognition in vitro by, 63
—, surface of, immunoglobulins on, 156, 1188
— with detectable surface-bound immunoglobulin, bone marrow as main source of, 1188
Lymphoid cell populations, immune, cell-to-cell interaction in, model to account for phenomenon of antigenic competition based on deficiency of, 1043
— cells, immune populations of, cellular deficit in reconstitutive capacity of, demonstrable in studies of delayed hypersensitivity in mice: evidence for thymus–bone marrow cell synergism, 1061
Lymphoma cells, nonsecretory, immunoglobulin in, synthesis and intracellular transport of, 901
Lymphotoxic and mitostatic effects of immunosuppressants and cytostatics, experimental system for simultaneous estimation of, 640
Lysin, Group C streptococcal phage-associated, purification and physical properties of, 1105

MACROPHAGE bactericidal capacity, immunologically mediated enhancement of, 1377
— DNA synthesis, activation of. Studies with inhibitors of RNA synthesis, 321
Macrophage functions, alterations of, by mediators from lymphocytes, 1356
— plasma membranes, 785, 807
Macrophage-digested antigen as inducer of delayed hypersensitivity, 494
Macrophage–melanocyte heterokaryons, 321
Macrophages, binding of immune complexes to, mechanisms underlying, 589
—, Macrophages, enzymatic degradation by, of hemoglobin to bile pigments, 1264
—, mouse peritoneal, and L-cells, interaction in vitro of *Mycoplasma pulmonis* with, 231
—, peritoneal, Syrian hamster, cytotoxic effects of mineral dusts on, 1149
—, restriction by, of herpes simplex virus, 19
Man, delayed-hypersensitive human donors, blood of, antigen-sensitive lymphocytes in, number and nature of, 740
—, dimeric IgA of human origin, ultrastructure of, further studies on, 1035
—, human fibroblast cultures, induction of foci and antigen in by filtrates; foci of altered cells and a common antigen, 454
—, human HL-A haplotypes, immunogenicity of, as measured by skin graft survival times and mixed leukocyte reactions, 411
—, malignant human plasmacytes and lymphocytes, synthesis and assembly of immunoglobulins by, 1118
—, placenta of, immunohistological and elution studies of, 1
—, pulmonary methane excretion in, factors influencing, 572
Melanocyte–macrophage heterokaryons, 321
Membrane lesions, ultrastructural,

production of by fifth component of complement, 53
Membranes, macrophage plasma, 785, 807
Membranolytic effects, differential, of microcrystalline sodium urate and calcium pyrophosphate dihydrate, 100
Memory, immunological, in vitro, 846
—, suppression of, by passive immunization late in primary response, 857
Metabolic activity and organ weights, effects of early environmental stresses on, 147
Metabolism, *in situ*, of methane-producing colonic bacteria, indirect method of studying, 572
Methane excretion, pulmonary, in man, factors influencing, 572
Methane-producing colonic bacteria, *in situ* metabolism of, indirect method of studying, 572
Microflora, mouse intestinal, with emphasis on strict anaerobes, 339
Mineral dusts, cytotoxic effects of, on Syrian hamster peritoneal macrophages, 1149
Mitostatic and lymphotoxic effects of immunosuppressants and cytostatics, experimental system for simultaneous estimation of, 640
Mononuclear cells, nature and specificity of, in experimental autoimmune inflammations and mechanisms leading to their accumulation, 1242
Mousepox: generalized viral infection–mechanisms of recovery from, 1074, 1090
Mycoplasma pulmonis, interaction in vitro with mouse peritoneal macrophages and L-cells, 231
Myocardial infarcts of rats, phlogistic role of C3 leukotactic fragments in, 885

NON–H-2 and H-2 antibody, differential immunofluorescence of fertilized mouse eggs with, 1282

OLIGOLYSINE antibody, anti–dinitrophenyl-, carrier specificity and genetic control of. Antigen recognition and antibody specificity, 1199
Organ weights, and metabolic activity, effects of early environmental stresses on, 147

PATHOGENESIS of H-1 virus infection of embryonic hamster bone in organ culture, 506
Pedicles of skin, alymphatic, allografts transplanted to, sensitization of rats by, 951
Peritoneal cavity, inflamed, lymphocyte traffic from blood into, 864
— cells, rat, IgE and IgGa antibody-mediated release of histamine from, 752, 772
— exudate cells, action on, 649
— macrophages, mouse, and L-cells, interaction in vitro of *Mycoplasma pulmonis* with, 231
— macrophages, Syrian hamster, cytotoxic effects of mineral dusts on, 1149
PHA stimulation, and alloantigen stimulation, of whole and gradient-separated subpopulations of spleen cells, effects of neonatal thymectomy and thymus graft restoration on, 63
Phage-associated lysin, Group C streptococcal, purification and physical properties of, 1105
(Phe, G)–Pro– –L, a synthetic polypeptide derived from multichain polyproline, immunocompetent precursors specific for two distinct regions within, frequency of, in inbred mouse strains, 216
Phlogistic role of C3 leukotactic fragments in myocardial infarcts of rats, 885

Placenta, human, immunohistological and elution studies of, 1
Plasma membranes, macrophage, 785, 807
Plasmacytes and lymphocytes, malignant human, synthesis and assembly of immunoglobulins by, 1118
Plasmin, digestion with, derivation of activators of prekallikrein from active Hageman factor by, 696
Platelets, leukocyte-dependent release of vasoactive amines from, and complement, role of, 554
Polynucleotides, synthetic, regulation by, of the immune system, 649, 665
Polypeptide, synthetic, (Phe, G)–Pro– –L, derived from multichain polyproline, frequency of immunocompetent precursors specific for two distinct regions within, in inbred mouse strains, 216
Polypeptides, synthetic, immune responses to, cellular basis of genetic control of, 216
Polyproline, multichain, (Phe, G)–Pro– –L, a synthetic polypeptide derived from, frequency of immunocompetent precursors specific for two distinct regions within, in inbred mouse strains, 216
Polysaccharide, *Escherichia coli*, antigen-binding cells in mice immune or tolerant to, 1015
Proliferative response, tempo and specificity of, and number of reactive cells from immunized donors, 442
Pulmonary methane excretion in man, factors influencing, 572
— tuberculosis, resistance to, temporal development of, in Swiss albino mice, 376
Pyrophosphate, calcium, dihydrate, and microcrystalline sodium urate, differential membranolytic effects of, 100

RADIATION, and cyclosphosphamide, contrasting effects of, on immune responses of mouse, 877
Reconstitutive capacity, cellular deficit in, of immune populations of lymphoid cells demonstrable in studies of delayed hypersensitivity in mice: evidence for thymus–bone marrow cell synergism, 1061
Rejection of tumor allografts by mouse spleen cells sensitized in vitro, 821
Renin, and angiotensin, in rat and dog brain, 353
Response, primary, suppression of memory by passive immunization late in, 857
Responses, humoral and cellular immune, to glucagon, specificity of: functional dissection of antigen molecule, 1294
Ribonucleic acid synthesis, studies with inhibitors of. Activation of macrophage DNA synthesis, 321
Rosette-forming cells, ^{3}H-thymidine–labeled, appearance of, 305

SARCOMA cells, murine, antigenic specificities on. Reciprocal relationship between normal transplantation antigens (H-2) and tumor-specific immunogenicity, 1171
Sensitization, kinetics of lymphocyte stimulation during, and recovery from tolerance, 187
Serum, heterologous anti-lymphocytic, immunosuppressive and graft-rejecting antibodies in, 81
Skin collagen, rat, distribution of antigenic determinants in, 1309
— graft survival times and mixed leukocyte reactions, immunogenicity of human HL-A haplotypes as measured by, 411
— homografts: tolerogenic vs. im-

munogenic influences in mice, 921
Skin, pedicles of, alymphatic, allografts transplanted to, sensitization of rats by, 951
Sodium urate, microcrystalline, and calcium pyrophosphate dihydrate, differential membranolytic effects of, 100
Spleen cells, mouse, sensitized in vitro, rejection of tumor allografts by, 821
— cells, whole and gradient-separated subpopulations of, effects of neonatal thymectomy and thymus graft restoration on alloantigen and PHA stimulation of, 63
Stimulation, lymphocyte, in cutaneous basophil hypersensitivity. Role of adjuvant, 202
—, lymphocyte, kinetics of, during sensitization and recovery from tolerance, 187
Streptococcal carbohydrates, rabbit antibodies to, relationship between relative binding affinity and electrophoretic behavior of, 424
—, Group C, phage-associated lysin, purification and physical properties of, 1105
Streptococci, presence of Group A variant-like antigen in streptococci of other groups, with special reference to Group N, 479
Stresses, early environmental, effects of, on metabolic activity and organ weights, 147
Synergism, thymus–bone marrow cell, evidence for: cellular deficit in reconstitutive capacity of immune populations of lymphoid cells demonstrable in studies of delayed hypersensitivity in mice, 1061
Synthesis and intracellular transport of immunoglobulin in nonsecretory lymphoma cells, 901

TARGET cell, interaction at, of IgGa and IgE, 772
— cells, in vitro preparation of with antibody and challenge with antigen, optimum conditions for, 752
Thymectomy, neonatal, effects of, and thymus graft restoration on alloantigen and PHA stimulation of whole and gradient-separated subpopulations of spleen cells, 63
Thymic origin, antigen-reactive cells of, action on, 665
Thymus graft restoration, and neonatal thymectomy, effects of, on alloantigen and PHA stimulation of whole and gradient-separated subpopulations of spleen cells, 63
Thymus–bone marrow cell synergism, evidence for: cellular deficit in reconstitutive capacity of immune populations of lymphoid cells demonstrable in studies of delayed hypersensitivity in mice, 1061
Thymus-derived cells, influence of during development of antibody-producing cells, 1325
Tissue, hamster cheek pouch, lymphatic status of, in relation to its properties as graft and graft site, 620
Tolerance, recovery from, sensitization and, kinetics of lymphocyte stimulation during, 187
Tolerogenic vs. immunogenic influences of skin homografts in mice, 921
Transplantation antigens (H-2), normal, and tumor-specific immunogenicity, reciprocal relationship between. Antigenic specificities on murine sarcoma cells, 1171
Tritiated thymidine, ^{3}H-thymidine-labeled rosette-forming cells, appearance of, 305
Tuberculosis, experimental, measurement of resistance to, in albino mice. 362
—, pulmonary, resistance to, temporal development of, in Swiss albino mice, 376

Tumor allografts, rejection of, by mouse spleen cells sensitized in vitro, 821
— factor responsible for angiogenesis, isolation of, 275
Tumor-specific immunogenicity and normal transplantation antigens (H-2), reciprocal relationship between. Antigenic specificities on murine sarcoma cells, 1171

ULTRASTRUCTURAL membrane lesions, production of, by fifth component of complement, 53
Urate, sodium, microcrystalline, and calcium pyrophosphate dihydrate, differential membranolytic effects of, 100

VIRAL genome, chromosomal integration of, implications regarding; G_{IX} system: cell surface allo-antigen associated with murine leukemia virus, 1334
Viral infection, generalized: mousepox–mechanisms of recovery from, 1074, 1090
Virus, Gross leukemia, wild-type, and pathogenesis of glomerulonephritis of New Zealand mice, 113
—, H-1, infection of embryonic hamster bone in organ culture, pathogenesis of, 506
—, herpes simplex, restriction of by macrophages, 19
—, murine leukemia, cell surface allo-antigen (G_{IX} system) associated with: implications regarding chromosomal integration of viral genome, 1334
Virus–cell interaction, analysis of, in macrophage restriction of herpes simplex virus, 19
Viruses, murine leukemia, major genetic locus affecting resistance to infection with, 1219, 1234